arXiv:hep-th/0610188v3 19 Jan 2007

Cosmologies with General Non-Canonical Scalar Field

†Wei Fang[1], ‡H.Q.Lu[1], Z.G.Huang[2]
[1]Department of Physics, Shanghai University,
Shanghai, 200444, P.R.China
[2]Department of Mathematics and Physics,
Huaihai Institute of Technology, Lianyungang, 222005, P.R.China

Abstract

We generally investigate the scalar field model with the lagrangian $L = F(X) - V(\phi)$, which we call it *General Non-Canonical Scalar Field Model*. We find that it is a special square potential(with a negative minimum) that drives the linear field solution($\phi = \phi_0 t$) while in K-essence model(with the lagrangian $L = -V(\phi)F(X)$) the potential should be taken as an inverse square form. Hence their cosmological evolution are totally different. We further find that this linear field solutions are highly degenerate, and their cosmological evolutions are actually equivalent to the divergent model where its sound speed diverges. We also study the stability of the linear field solution. With a simple form of $F(X) = 1 - \sqrt{1-2X}$ we indicate that our model may be considered as a unified model of dark matter and dark energy. Finally we study the case when the baryotropic index γ is constant. It shows that, unlike the K-essence, the detailed form of $F(X)$ depends on the potential $V(\phi)$. We analyze the stability of this constant γ_0 solution and find that they are stable for $\gamma_0 \leq 1$. Finally we simply consider the constant c_s^2 case and get an exact solution for $F(X)$.

Keywords: Non-Canonical Scalar Field; Linear field solution; Dark Energy; K-essence; Cosmology.

PACS: 98.80.Cq, 04.65.+e, 11.25.-w

† xiaoweifang_ren@shu.edu.cn
‡ Alberthq_lu@staff.shu.edu.cn

1 Introduction

Dark energy problem may be one of the biggest issues in current theoretical physics and cosmology(see Ref[1] for a recent review). The building of theoretical models as well as the breakthrough at astrophysical observations have never halted since 1998. Technically speaking, one can operate on either the r.h.s or the l.h.s of the Einstein equation to get a reasonable interpretation of the accelerating expansion of the universe. Many candidates of dark energy have been proposed, such as the cosmological constant[2], quintessence[3], K-essence[4], phantom[5], modifying gravity[6] and so on. Among these models the scalar field models are undoubtedly the most important class of theoretical models. Generally speaking, the lagrangian of the scalar field model can be generally represented as[7]

$$L = f(\phi)F(X) - V(\phi) \tag{1}$$

where $X = \frac{1}{2}\nabla_\mu\phi\nabla^\mu\phi = \frac{1}{2}\dot{\phi}^2$ for a spatially homogeneous scalar field. Eq.(1) has included all the popular single scalar field models. It describes K-essence when $V(\phi) = 0$ and standard quintessence when $f(\phi)$ =constant and $F(X) = X$. The idea of K-essence was firstly introduced as a possible model for inflation[8] and Later was considered as a possible model for dark energy[4,9]. L.P.Chimento found the first integral of the K-essence field equation for any function $F(X)$ when the potential is taken as inverse square form or a constant[10]. In Ref[11], it was found that every quintessence model can be view as a K-essence model generated by a kinetic linear $F(X)$ function, and some K-essence potentials and their quintessence correspondence are also found.

In this paper, we will focus on another class of models with its lagrangian $L = F(X) - V(\phi)$, which we think is as important as K-essence model but whose general characters and roles in cosmology is far beyond clear. Here we should point out that, from the original literature's point of view[8,9], our lagrangian(Eq.(2)) also belongs to K-essence model, which is charactered by a lagrangian $L = L(X, \phi)$. However as far as we know, most works about K-essence model are just based on the lagrangian form $L = V(\phi)F(X)$. Therefore in this paper we call the model with lagrangian $L = F(X) - V(\phi)$ *Non-Canonical Scalar Field Model* while specify the model with the lagrangian $L = V(\phi)F(X)$ as K-essence model. The paper is organized as follows: Section 2 is the theoretical framework. In section 3 we investigate the linear field solution and find the potential driving this linear field solution. The divergent model with its speed of sound diverge($c_s^2 \to \infty$) is considered in section 4. Some solvable general non-canonical scalar field solution are discussed in section 5. Section 6 is the conclusion.

2 Basic Framework

Let us restrict ourselves for the time being to the cosmological setting corresponding to the flat universe described by the FRW metric. We consider the spatially homogeneous real scalar field ϕ with a non-canonical kinetic energy term. The lagrangian density is

$$L = F(X) - V(\phi) \tag{2}$$

where $V(\phi)$ is a potential and $F(X)$ is an arbitrary function of X. Obviously above equation is a special case of Eq.(1) when the function $f(\phi)$ =constant. It includes quintessence $[F(X) = X]$ and a phantom field $[F(X) = -X]$. In fact this form of lagrangian has appeared in Refs[1,12].

We can easily get the following equations:

$$p = L = F(X) - V(\phi) \tag{3}$$

$$\rho = 3H^2 = 2L_{,X}X - L = 2XF_{,X} - F(X) + V(\phi) \tag{4}$$

$${c_s}^2 = p_{,X}/\rho_{,X} = [1 + 2X\frac{F_{,XX}}{F_{,X}}]^{-1} \tag{5}$$

Where we take $8\pi G = 1$ for convenience. From Eqs.(3,4) we get the relation $\rho+p = 2XF_{,X}$ and this yields the state equation ω_ϕ larger than -1 if $F_{,X} > 0(\omega_\phi < -1$ if $F_{,X} < 0)$. Vikman has argued that it is impossible for ω_ϕ to cross the phantom line divide$(\omega_\phi = -1)$ in single scalar field theory[13]. However it is argued that this result holds only for models without considering higher derivative terms[14]. Eq.(5) describes the effective sound speed of the perturbations. ${c_s}^2 \equiv 1$ if $F_{,XX} = 0$ is satisfied.

The motion equations of the general non-linear scalar field are

$$\ddot{\phi} + 3{c_s}^2 H\dot{\phi} + \frac{\rho_{,\phi}}{\rho_{,X}} = 0 \tag{6}$$

$$(F_{,X} + 2XF_{,XX})\ddot{\phi} + 3HF_{,X}\dot{\phi} + V_{,\phi} = 0 \tag{7}$$

$$(\frac{\gamma}{\dot{\phi}})^{\cdot} + 3H(1-\gamma)(\frac{\gamma}{\dot{\phi}}) + \frac{V_{,\phi}}{3H^2} = 0 \tag{8}$$

where "$f_{,x}$" denotes the derivative with respect to subscript index x and $\gamma = (\rho + p)/\rho$ is the baryotropic index. Eqs.(6,7,8) are different forms of the motion equation and they are equivalent to each other.

3 The Linear Field Model

In this section we will investigate a special case that the field possesses a linear field solution $\phi = \phi_0 t$. We find that the form of potential $V(\phi)$ which drives this evolution

is a square potential with a negative minimum. We also show that the usual linear field solution in non-linear scalar field theory leads to an entirely different universe comparing with the universe in K-essence model.

For the linear field solution, $X = \frac{1}{2}\dot{\phi}^2 \equiv \frac{1}{2}{\phi_0}^2$ is a constant and $\ddot{\phi} = 0$, then we get following equation from Eqs.(4,7):

$$\frac{{V_{,\phi}}^2}{3{F_{,X}}^2{\phi_0}^2} - V(\phi) + F(X) - {\phi_0}^2 F_{,X} = 0 \tag{9}$$

F and $F_{,X}$ are only the function of X and therefore they both are constant. We set $F_0 = F(X_0)$ and $F_{,0} = F_{,X}(X_0)$ by evaluating them at $X = X_0 = \frac{1}{2}{\phi_0}^2$. Solving Eq.(9) and Eq.(4) we get the exact solutions for potential $V(\phi)$ and scale factor a:

$$V(\phi) = \frac{3}{4}{F_{,0}}^2{\phi_0}^2(\phi + c)^2 + F_0 - F_{,0}{\phi_0}^2 \tag{10}$$

$$a = a_0 exp[-\frac{F_{,0}}{4}(\phi_0 t + c)^2] \tag{11}$$

Therefore the linear field solution leads to a square potential Eq.(10) and a evolution of scale factor Eq.(11). It is worthwhile to compare the same case in K-essence model. It is argued[15] that the same linear field solution in K-essence model leads to an inverse square potential and a power law expansion of scale factor. It is interesting that the same linear field solution lead to different cosmological evolution and therefore different cosmological implication. We should emphasize that the potential is exactly derived from Eqs.(4,9) and its form is unique. Moreover in this case the different forms of $F(X)$ degenerate to only two cases: $F_{,0} > 0$ and $F_{,0} < 0$ and respectively correspond to non-phantom($\omega > -1$) and phantom case($\omega < -1$). The phantom case($F_{,0} < 0$) describes a universe from contracting phase to expanding phase and is excluded easily by current observation. So we restrict $F_{,0} > 0$ for next discussion. From Eq.(4) we get $\rho = F_{,0}{\phi_0}^2 - F_0 + V(\phi)$. To ensure the energy density ρ has a positive kinetic energy term we demand $F_{,0}{\phi_0}^2 - F_0 > 0$ and this immediately leads to the square potential with a negative minimum value $F_0 - F_{,0}{\phi_0}^2$[see Eq.(10)]. In fact this result is well-intelligible in an expanding universe. Because if the square potential has a non-negative minimum it is well-known that the scalar field ϕ will roll down the potential and finally cease at the minimum position $\phi = -c$ and the linear field solution $\phi = \phi_0 t$ will be no long valid. On the other hand it is argued that for a potential with a negative minimum the scalar field can oddly roll up the potential from its minimum(see Fig.1) and the universe enters a contracting phase from an expanding phase[16] and therefore the scalar field can evolve to ∞.

It is very interesting that the universe in our model can avoid a beginning singularity. If we think the classic cosmology is valid when energy scale is below Plank scale, then the scale factor at the beginning is very small[$a = a_0 e^{-\rho_{pl}/(3F_{,0}{\phi_0}^2)}$, where ρ_{pl} is the energy

density at plank time] but does not equal zero. However this universe can not escape from a collapse in future. This evolutive behaviors are completely different from the same linear field solution case($\phi = \phi_0 t$) in K-essence model where the scale factor behaves as $a \propto t^n$[10] and the universe was birth from a singularity and expand for ever.

One of our concerns is whether our model can describe a suitable universe with a phase of accelerating expansion. The answer is positive because we have $\ddot{a} \propto -(\rho + 3p) = -\frac{3}{2}F_{,0}{\phi_0}^2[2 - F_{,0}(\phi + c)^2]$ and $\rho + 3p < 0$ for $\phi < \phi_1$ or $\phi > \phi_2$, where $\phi_1 = -c - \sqrt{\frac{2}{F_{,0}}}, \phi_2 = -c + \sqrt{\frac{2}{F_{,0}}}$. The potential and the evolutive behaviors of universe are showed in Fig.1. We can see from Fig.1, the field rolls down the potential from an initial value and the universe undergoes an accelerating expansion. When the field evolutes to ϕ_1, the universe enters a decelerating expansion and finally becomes zero expansion rate when the field arrives at $\phi = -c$. When the field crosses the point $\phi = -c$ the expansion rate H dramatically changes it sign from $H > 0$ to $H < 0$ and the field rolls up the potential from its minimum. When the field rolls from $-c$ to ϕ_2 the universe undergoes an accelerating contraction. After going over ϕ_2, the universe enters a decelerating contraction and collapses to singularity finally.



In addition, since we get the potential(Eq.(10)) just from the linear field solution without any other assumption, another non-trivial question is whether an arbitrary square potential with a negative constant can always lead to the linear field solution. Unfortunately, we will demonstrate that it is not the case: not all the square potential with a negative constant can possess the linear field solution.

Given a arbitrary potential:

$$V(\phi) = A(\phi + c)^2 - B \tag{12}$$

where A, B are arbitrary positive constants. In order to have a linear field solution, from Eq.(10) A and B should satisfy $A = \frac{3}{4}{F_{,0}}^2{\phi_0}^2, B = F_0 - F_{,0}{\phi_0}^2$. So we get the following

constraint equation:

$$4A = 3F_{,0}(F_0 - B) \tag{13}$$

It means that A and B are not arbitrary parameters. In other words, given a value of A, the value of B is determined by Eq.(13). i.e, only the square potential with its parameters satisfying Eq.(13) can lead to a linear field solution. From Eq.(13) we find another interesting character for the linear field solution. Since there only appears the first two coefficients $(F_0, F_{,0})$ of the series expansion of the function $F(X)$ around $X = X_0$(we can expand function $F(X)$ as $F_0 + F_{,0}(X - X_0) + \cdots$ around X_0). The evolution of universe will be thought of as equivalent as long as the function $F(X)$ has the same first two coefficients of its expansion, disregarding the rest of the higher order terms. This means that the linear filed solution model possesses a high degenerate character.

The last question we ask is, how stable are the linear field solutions. Since for the linear field solution we have $3HF_{,0}\phi_0 + V_{,\phi} = 0$, we can rewrite Eq.(7) as follows:

$$\frac{d\dot{\phi}}{dt} = \frac{-3H}{F_{,X} + 2XF_{,XX}}(F_{,X}\dot{\phi} - F_{,0}\phi_0) \tag{14}$$

For the case that $c_s^2 = costant \geq 0$, Eq.(14) becomes

$$\frac{d\dot{\phi}}{dt} = -3Hc_s^2(\dot{\phi} - \phi_0) \tag{15}$$

Integrating Eq.(15), we have

$$\dot{\phi} = \phi_0 + \frac{c_1}{a^{3c_s^2}} \tag{16}$$

Eq.(16) shows that, if the universe expands for ever, $\dot{\phi}$ has an asymptotic limit ϕ_0 and the solution $\phi = \phi_0 t$ is stable. However, as we known from Eq.(11), after expanding to a maximum scale factor the universe will contract to a singularity eventually. Therefore this linear field solution can not be stable.

4 The Divergent Model

In this section we investigate another interesting model where the speed of sound is divergent($c_s^2 \to \infty$). From Eq.(5), we have:

$$1 + 2X\frac{F_{,XX}}{F_{,X}} = 0 \tag{17}$$

Integrating Eq.(17) we get the form of function $F(X)$:

$$F(X) = c_2 X^{\frac{1}{2}} + c_3 \tag{18}$$

where c_2, c_3 are the integral constants. From Eq.(18), the sound speed of this special form of $F(X)$ diverges(${c_s}^2 \to \infty$). The same form of function $F(X)$ is also obtained in K-essence model with the lagrangian being $-V(\phi)F(X)$[10,15]. Recently This type lagrangian is thoroughly investigated and exploited in model building[17], where this type lagrangian is called as *Cuscuton*. From Eq.(4) and Eq.(18), we have

$$3H^2 = V(\phi) - c_3 \tag{19}$$

From Eq.(7) and Eq.(19) we get

$$\frac{1}{2}c_2\sqrt{3(V(\phi) - c_3)} + dV(\phi)/d\phi = 0 \tag{20}$$

Therefore we can immediately get the potential $V(\phi)$:

$$V(\phi) = \frac{3}{8}c_2^2(\phi - c_4)^2 + c_3 \tag{21}$$

where c_4 is an integral constant. For the model building the concrete form of function $F(X)$ and the potential $V(\phi)$ can be constructed respectively, however, it is very interesting that the divergent model with the special lagrangian Eq.(18) determine the unique form of potential Eq.(21). This maybe means that the divergent model with the speed of sound $c_s^2 = \infty$ has some special implications. Let us recall the result obtained in Section 3: if the square potential satisfies the constraint Eq.(10), the solution of the scalar field will be linear field solution. From Eqs.(10, 18), we have $\frac{3}{4}{F_{,0}}^2{\phi_0}^2 = \frac{3}{8}c_2^2, F_0 - F_{,0}{\phi_0}^2 = c_3$, which just coincides with Eq.(21). This means that the divergent model and the linear field solution are degenerate. Namely the divergent model and the linear field solution model are kinetically isomorphic and share the same evolution of scale factor a and scalar field ϕ:

$$\phi = \phi_0 t, \quad a = a_0 exp[-\frac{F_{,0}}{4}(\phi_0 t + c)^2] \tag{22}$$

However it remains to clarify why this could happen. From the mathematical point of view, we maybe get the interpretation from the motion equation of field Eq.(7) since both the divergent model and the linear field model lead the first term of Eq.(7) to vanish. Therefore they share the same motion equation and then lead to the same cosmological evolution. But what is the physical implication that the scalar field theory with an infinite speed of sound is degenerate with the linear field model? There also exists the same situation in the K-essence model that the linear field solution and the divergent model($c_s^2 = \infty$) are isomorphic[18].

5 Solvable General Non-canonical Scalar Field Cosmologies

In this section we will focus on some special cases when Eq.(8) exists a first integral or can be solved exactly. Though what we consider are quite simple, they can also lead to some important results. Additional, we will try to find the relationships and the differences between our General Non-Canonical Scalar Field model and K-essence model.

A. $V(\phi) = V_0$

When the potential $V(\phi)$ is a constant($= V_0$), Eq.(8) exists a first integral

$$\frac{\gamma}{\dot{\phi}} = \frac{c_5}{a^3 H^2} \tag{23}$$

where c_5 is an arbitrary integral constant. Eq.(23) is very similar with the first integral obtained in K-essence model with a constant potential[10,19]. For a constant potential the K-essence lagrangian can be written as $L_k = -V_0 F_k(X)$ while the lagrangian in our model is $L_g = F_g(X) - V_0$. They are actually equivalent if we define $F_g(X) = V_0(1 - F_g(X))$. So our model can easily reproduce the K-essence models with constant potential. The K-essence models with a constant potential are hotly studied for its exquisite role in unifying the dark matter and dark energy[20]. Let we consider a special form of lagrangian $L_g = 1 - \sqrt{1 - 2X} - V(\phi)$, which is considered as a Non-Linear Born-Infeld(NLBI) scalar field theory in Ref[21]. When the potential is constant V_0, we can find the exact solutions:

$$\dot{\phi}^2 = \frac{c_6}{c_6 + a^6} \tag{24}$$

$$\rho = \sqrt{1 + \frac{c_6}{a^6}} + (V_0 - 1) \tag{25}$$

$${c_s}^2 = 1 - \dot{\phi}^2 = \frac{a^6}{c_6 + a^6} \tag{26}$$

Where c_6 is an integral constant. From Eqs.(25,26) we know that, the energy density behaves as dark matter at early time(for small a, $\rho \propto a^{-3}$ and ${c_s}^2 \simeq 0$) and dark energy at late time(for large a, $\rho \simeq V_0 = const$ and ${c_s}^2 \simeq 1$). So, our model can also play the same role that unifies the dark matter and dark energy.

B. $\gamma =$**constant**

In this subsection we will assume that the baryotropic index $\gamma = \gamma_0 =$constant. Then the state equation $\omega = \gamma - 1 = \gamma_0 - 1$, is also a constant. The constant γ kinematically leads to the cosmological solution

$$a = a_0 t^{2/3\gamma_0}, \quad \rho_\phi = \frac{4 a_0^{3\gamma_0}}{3\gamma_0^2} / a^{3\gamma_0} \tag{27}$$

From the relation

$$\gamma_0 = \frac{\rho + p}{\rho} = -\frac{2\dot{H}}{3H^2} = \frac{2XF_{,X}}{2XF_{,X} - F(X) + V(\phi)} \tag{28}$$

We get the following equation:

$$\frac{2(\gamma_0 - 1)}{\gamma_0} XF_{,X} - F(X) + V(\phi) = 0 \tag{29}$$

In Ref[18], it is showed that $F(X) = X^{\frac{\gamma_0}{2(\gamma_0 - 1)}}$ can lead to the constant γ with any potential in K-essence model. Here we show that in our model the form of $F(X)$ depends on the potential(see Eq.(29)). Namely, to admit the cosmological solution Eq.(27), the kinetic term $F(X)$ and potential term $V(\phi)$ must satisfy Eq.(29). Only for a constant potential V_0 ,we get a similar function of $F(X)$:

$$F(X) = c_7 X^{\frac{\gamma_0}{2(\gamma_0 - 1)}} + V_0 \tag{30}$$

where c_7 is an integral constant.

It is quite interesting to consider whether the solution with constant γ_0 is stable. To answer this question, we let γ vary with time. Differentiating the equation of the baryotropic index γ, we get

$$\dot{\gamma} = (\gamma - 1)(3H\gamma + \frac{\dot{p}}{p}) \tag{31}$$

We immediately get two critical points: $\gamma_0 - 1 = 0$ or γ_0 satisfies Eq.(32):

$$3H\gamma_0 + \frac{\dot{p}}{p} = 0 \tag{32}$$

When this stationary condition Eq.(32) holds, the potential $V(\phi)$ and function $F(X)$ will satisfy the relation:

$$p = F(X) - V(\phi) = \frac{c_8}{a^{3\gamma_0}} \tag{33}$$

From Eqs.(31,32), we have

$$\dot{\gamma} = 3H(\gamma - 1)(\gamma - \gamma_0) \tag{34}$$

Integrating Eq(34) we get

$$\gamma = \frac{\gamma_0 a^{3(1-\gamma_0)} - c_9}{a^{3(1-\gamma_0)} - c_9} \tag{35}$$

where c_8, c_9 is an integral constant. Eq.(35) indicates that, for the expanding universe and $\gamma_0 < 1$ the baryotropic index γ has the asymptotic limit γ_0. However for $\gamma_0 > 1$ the baryotropic index γ will approach the asymptotic limit 1. The case $\gamma_0 = 1$ should be considered apart. For $\gamma_0 = 1$, the solution is

$$\gamma = 1 - \frac{c_{10}}{3lna} \tag{36}$$

Where c_{10} is an integral constant. Eq.(36) shows that the solution with $\gamma_0 = 1$ is also stable in an expanding universe. Therefore, we can conclude that the solutions with constant baryotropic index are attractors in the case $\gamma_0 \leq 1$ and the $\gamma_0 = 1$ solutions separate stable from unstable regions in the phase space.

C. c_s^2 **=constant**

For the dark energy behaving as a fluid, the speed of sound c_s is another important parameter in addition to the equation of state ω. The speed of sound c_s is the propagation of the perturbation of the background scalar field, which can affects the CMB power spectrum. Therefore the effective sound speed c_s of dark energy would provide crucial information which is complementary to the equation of state ω. In this subsection we will study the simple case that the speed of sound is constant. From Eq.(5), we obtain the equation as follows:

$$2c_s^2 X F_{,XX} = (1 - c_s^2) F_{,X} \tag{37}$$

Integrating Eq.(26), we have

$$F(X) = \frac{2c_s^2}{1 + c_s^2} c_{11} X^{\frac{1+c_s^2}{2c_s^2}} + c_{12} \tag{38}$$

where $c_{11},\ c_{12}$ is the integral constants.

6 Conclusion

In this paper we have generally investigated the general non-canonical scalar field model as a candidate of dark energy. We found that it was a special square potential(with a negative minimum) that drove the linear field solution in our model while the potential should be taken as an inverse square form in K-essence model. Our results showed that the linear field solution was highly degenerate, and shared the same cosmological evolution with the divergent model where its sound speed $c_s^2 \to \infty$. We pointed out that our model with a constant potential was actually equivalent to the K-essence model also with a constant potential. With a simple form of $F(X)$ we indicated that our model can be also considered as a unified model of dark matter and dark energy. In addition we studied the constant γ_0 case. The results showed that, unlike the K-essence model, the detailed form of $F(X)$ depended on the potential. We find that the constant γ_0 solution is stable for $\gamma_0 \leq 1$. We also found the form of $F(X)$ which possessed the constant c_s^2 solution. Our work may throw light on the study of the scalar field theory and the exploration of dark energy.

7 Acknowledgement

W.Fang would like to thank Dr.Hongwang Yu and Xingyu Jin for useful discussion. We also appreciate the anonymous referee for the helpful suggestion. This work is partly supported by NNSFC under Grant No.10573012 and No.10575068 and by Shanghai Municipal Science and Technology Commission No.04dz05905.

References

1. E.J.Copeland, M.Sami and S.Tsujikawa, hep-th/0603057

2. S.Weinberg, Rev.Mod.Phys**61**(1), 1(1989);
   P.J.Steinhardt, L.Wang and I.Zlatev,Phys.Rev.D**59**, 123504(1999);
   P.J.E.Peebles and B.Ratra, Rev.Mod.Phys**75**(2), 559(2003);
   A.G.Riess et al., Astrophys.J**607**, 665-687(2004);
   G.W.Gibbons, hep-th/0302199.

3. R.R.Caldwell, R.Dave and P.J.Steinhardt, Phys.Rev.Lett.**80**, 1582(1998);
   P,J.Steinhardt,L.Wang and I.Zlatev, Phys.Rev.Lett.**82**, 896(1996);
   X.Z.Li, J.G.Hao and D.J.Liu, Class.Quantum Grav.**19**, 6049(2002).

4. J.M.Aguirregabiria, L.P.Chiemento and R.Lazkoz, Phys.Rev.D**70**, 023509(2004);
   E.J.Copeland, M.R.Garousi, M.Sami and S.Tsujikawa, Phys.Rev.D**71**, 043003(2005);
   G.Panotopoulos; astro-ph/0606249;
   M.Malquarti, E.J.Copeland and A.R.Liddle, Phys.Rev.D**68**, 023512(2003);
   R.Lazkoz, Int.J.Mod.Phys.D**14**, 635-642 (2005).

5. R.R.Caldwell, Phys.Rev.Lett.B**545**, 23(2002);
   M.Sami, Mod.Phys.Lett.A**19**, 1509(2004);
   M.Sami and T.Padamanabhan, Phys.Rev.D**67**, 083509(2003);
   M.Sami, P.Chingangbam and T.Qureshi, hep-th/0301140;
   P.Singh, M.Sami, N.Dadhich,Phys.Rev.D**68**, 023522(2003);
   V.Faraoni, Class.Quant.Grav.**22**, 3235-3246(2005);
   D.J.Liu and X.Z.Li, Phys.Rev.D**68**, 067301(2003);
   M.P.Dabrowski et.al., Phys.Rev.D**68**, 103519(2003);
   S.M.Carroll, M.Hoffman and M.Teodden, Phys.Rev.D**68**, 023509(2003);
   Y.S.Piao, R.G.Cai, X.M.Zhang and Y.Z.Zhang, Phys.Rev.D**66**, 121301(2002);
   S.Mukohyama, Phys.Rev.D**66**, 024009(2002);
   T.Padmanabhan, Phys.Rev.D**66**, 021301(2002);
   G.Shiu and I.Wasserman, Phys.Lett.B**541**, 6(2002);
   L.kofman and A.Linde, JHEP**0207**, 004(2002);
   A.Sen, JHEP**0210**, 003(2002);

I.Ya.Aref'eva and A.S.Koshelev, hep-th/0605085;
I.Ya.Aref'eva, A.S.Koshelev and S.Yu.Vernov, Phys.Rev.D**72**, 064017(2005);
I.Ya.Aref'eva, A.S.Koshelev and S.Yu.Vernov, Phys.Lett.B**628**, 1-10(2005);
I.Ya.Aref'eva, A.S.Koshelev and S.Yu.Vernov, astro-ph/0412619;
N.Moeller and B.Zwiebach, JHEP**0210**, 034(2002);
P.Mukhopadhyay and A.Sen, JHEP**0211**, 047(2002);
T.Okunda and S.Sugimoto, Nucl.Phys.B**647**, 101-116(2002);
G.Gibbons, K.Hashimoto and P.Yi, hep-th/0209034;
B.Chen, M.Li and F.Lin, hep-th/0209222;
G.Felder, L.Kofman and A.Starobinsky, JHEP**0209**, 026(2002);
M.C.Bento, O.Bertolami and A.A.Sen, hep-th/0208124;
H.Lee et.al., hep-th/0210221;
J.G.Hao and X.Z.Li, Phys.Rev.D**66**, 087301(2002); Phys.Rev.D**68**, 043501(2003);
X.Z.Li and X.H.Zhai, Phys.Rev.D**67**, 067501(2003);
Z.G.Huang, H.Q.Lu and W.Fang, Class.Quant.Grav**23**, 6215(2006);
S.Nojiri and S.D.Odintsov, Phys.Lett.B**562**, 147(2003);
E.Elizalde, S.Nojiri and S.D.Odintsov, Phys.Rev.D**70**, 043539(2004);
S.Nojiri and S.D.Odintsov, Phys.Rev.D**70**, 103522(2004);
S.Nojiri, S.D.Odintsov and S.Tsujikawa, Phys.Rev.D**71**,063004(2005);
S.Nojiri and S.D.Odintsov, Phys.Rev.D**72**, 023003(2005).

6. G.Dvali, G.Gabadadze and M.Porrati, Phys.Lett.B**485**, 208(2000);
G.Dvali, G.Gabadadze, M.Kolanovic and F.Nitti, Phys.Rev.D**64**, 084004(2001);
G.Dvali, G.Gabadadze, M.Kolanovic and F.Nitti, Phys.Rev.D**65**, 024031(2002);
C.Deffayet, Phys.Lett.B**502**, 199(2001);
C.Deffayet, G.R.Dvali and G.Gabadadze, Phys.Rev.D**65**, 044023(2002);
S.Nojiri and S.D.Odintsov, hep-th/0601213;
B.Li and M.C.Chu, Phys.Rev.D**74**, 104010(2006);
B.Li, K.C.Chan and M.C.Chu, astro-ph/0610794.

7. A.Melchiorri, L.Mersini, C.J.Odman and M.Trodden, Phys.Rev.D**68**, 043509(2003).

8. J.Garriga and V.F.Mukhanov, Phys.Lett.B**458**, 219(1999);
C.Armendáriz-Picón, T.Damour and V.Mukhanov, Phys.Lett.B**458**, 209(1999).

9. C.Armendáriz-Picón, V.Mukhanov and P.J.Steinhardt, Phys.Rev.Lett**85**, 4438(2000);
C.Armendáriz-Picón, V.Mukhanov and P.J.Steinhardt, Phys.Rev.D**63**, 103510(2001);
T.Chiba, Phys.Rev.D**66**, 063514;
T.Chiba, T.Okabe and M.Yamaguchi, Phys.Rev.D**62**, 023511(2000);
M.Malquarti, E.J.Copeland, A.R.Liddle and M.Trodden, Phys.Rev.D**67**, 123503(2003).

10. L.P.Chimento, Phys.Rev.D**69**, 123517(2004).

11. J.M.Aguirregabiria, L.P.Chimento and R.Lazkoz, Phys.Lett.B**631**, 93-99(2005).

12. V.Mukhanov and A.Vikman, JCAP**0602**004 (2006); A.Vikman, astro-ph/0606033
E.Babichev, hep-th/0608071.

13. A.Vikman, Phys.Rev.D**71**, 023515(2005).

14. M.Li, B.Feng and X.Zhang, JCAP**0512**,002(2005);
A.Anisimov, E.Babichev and A.Vikman, JCAP**0506**,006(2005);
X.F.Zhang and T.Qiu, astro-ph/0603824.

15. L.P.Chimento and A.Feinstein, Mod.Phys.Lett.A**19**, 761-768 (2004).

16. G.Felder, A.Frolov, L.Kofman and A.Linde, Phys.Rev.D**42**, 023507(2002);
A.de la Macorra and C.Stephan, Phys.Rev. D**65**, 083520(2002);
Imogen P. C. Heard and David Wands, Class.Quant.Grav.**19**, 5435-5448(2002);
L.Perivolaropoulos, Phys.Rev. D**71**, 063503(2005);
A.de la Macorra and G. German, astro-ph/0212148;
W.Fang, H.Q.Lu and Z.G.Huang, hep-th/0606033.

17. N.Afshordi, D.J.H.Chung and G.Geshnizjani, hep-th/0609150.

18. L.P.Chimento and A.Feinstein, Mod.Phys.Lett.A**19**, 761-768 (2004).

19. C.Armendáriz-Picón,T.Damour and V.Mukhanov,Phys.Lett.B**458**, 209(1999).

20. R.J.Scherrer, Phys.Rev.Lett**93**, 011301(2004);
L.P.Chimento, M.Forte and R.Lazkoz, Mod.Phys.Lett.A**20**, 2075(2005);
M.Novello, M.Makler, L.S.Werneck and C.A.Romero, Phys.Rev.D**71**, 043515(2005)

21. H.Q.Lu, Int.J.Mod.Phys.D**14**, 355(2005);
W.Fang, H.Q.Lu, Z.G.Huang and K.F.Zhang, Int.J.Mod.Phys.D**15**, 199-214(2006)(hep-th/0409080);
W.Fang, H.Q.Lu and B.Li and K.F.Zhang, Int.J.Mod.Phys.D**15**, 1947-1961(2006)(hep-th/0512120);
Z.G.Huang, X.H.Li and Q.Q.Sun, hep-th/0610019